



TRIALS OF VITAMIN SUPPLEMENTS FOR THE PREVENTION OF CHRONIC DISEASE

Comment by Pamela Mason

Vitamins are essential for our health and wellbeing. A significant number of British people have diets lacking in these essential nutrients. Vitamins and minerals should ideally be obtained from the diet, but failing that a supplement can help to fill the nutritional gap. To expect a vitamin pill to prevent a multi-factorial disease which has taken a lifetime to develop, may be unrealistic, particularly if the person already has the disease in question. But what can be expected is that a multivitamin containing a wide variety of vitamins and minerals in the recommended daily amount will ensure that the person's intake does not fall below recommended levels.

During recent years, there have been a number of research studies evaluating the effects of vitamin and mineral supplements in the prevention of chronic disease such as cardiovascular disease, cancer, osteoporosis and eye diseases (eg, cataract and age-related macular degeneration).

Results from these studies have been inconsistent. Some studies have concluded that vitamin supplements are associated with reduced risk of chronic disease while others have not.

Cardiovascular disease

The Nurses' Health Study found that the risk of heart disease was reduced by 24% in those who regularly used multivitamins,¹ while in the Stockholm Heart Epidemiology Program (SHEEP), risk of myocardial infarction (MI) was significantly reduced in both men and women who took multivitamins.²

Folic acid (with or without vitamin B6 and B12) has been linked with reduced risk of cardiovascular disease by virtue of its ability to lower homocysteine, with recent meta-analyses confirming these findings.^{3,4} In the Cambridge Heart Antioxidant Study (CHAOS), vitamin E reduced the 1-year rate of non-fatal myocardial infarction.⁵

However, two large randomised controlled trials published in 2006 (the Heart Outcomes Prevention Evaluation⁷ and the NORVIT⁸ Trial) found that supplements of B12, folic acid and B6 in combination reduced plasma homocysteine, but did not reduce the risk of major cardiovascular events in patients with cardiovascular disease⁷.

Cancer

In the Health Professionals' Follow-up Study, men who reported folate consumption from multivitamins for less than 10 years had a 25% reduction in colon cancer risk.⁹ The Nurses' Health Study found that women who reported multivitamin use (with folate) of 15 years or more had a 75% reduction in colorectal cancer risk.¹⁰

An intervention trial in China¹¹ found combined daily doses of beta-carotene, vitamin E and selenium over 5 years were linked with a 13% reduction in cancer deaths and a 9% reduction in all-cause mortality. However, a recent observational study concluded that high use of vitamins and other supplements may increase the risk of advanced fatal prostate cancer.¹³ But, the results of this **study are at odds** with a recent case-control study that showed that increased blood levels of selenium in

combination with a daily multivitamin supplement reduced the risk of prostate cancer by 40%.¹⁴

What is going on?

Clearly no firm conclusions can be reached about the effects of supplements in the prevention of chronic disease such as cardio vascular disease and cancer. Research findings are conflicting and inconclusive.

What is clear, **however, is that vitamins and minerals are essential nutrients which are vital for health.** Furthermore, surveys have shown that the diet of many people in the UK is lacking in these essential substances. **Among women of childbearing age, nine in ten women fail to achieve an adequate intake of either folic acid or iron. Half of adult men and three quarters of adult women fail to achieve the Reference Nutrient Intake for magnesium. More worryingly still, one in ten men and one in seven women fail to achieve the Lower Reference Nutrient Intake (LRNI) an intake below which deficiency is likely, of magnesium.**¹⁵

Studies in adults have shown that **supplement use can make a significant contribution to vitamin and mineral intake.** The National Diet and Nutrition Survey (NDNS) in British adults found that supplement users had higher intakes of vitamins and minerals and were less likely to have intakes below the Reference Nutrient Intake (RNI) than non-supplement users.¹⁵ Similar findings have been shown in Ireland,¹⁶ Germany,^{17,18} the US¹⁹ and Canada.²⁰ In youngsters too, supplements have been shown to make a substantial contribution to the intakes of vitamins and minerals in both toddlers^{21,22} and teenagers.²³⁻²⁶ Several studies²⁷⁻²⁹ have also shown that supplementation with vitamin and minerals can improve plasma levels of micronutrients and reduce the prevalence of suboptimal plasma concentrations.

Clearly supplementation can help to bridge the nutritional gap observed in significant numbers of people, as well as maintaining good health.

When it comes to prevention of chronic disease in middle and older age it was never the intention to use vitamin supplements primarily for the prevention of chronic disease. Vitamins are intended primarily for health maintenance. Moreover, it would seem very optimistic to expect a vitamin pill to prevent heart disease or cancer. Vitamins will prevent or cure vitamin deficiency simply because there is one cause of deficiency – i.e. lack of the vitamin or mineral concerned.

The development of chronic conditions like cardiovascular disease and cancer is related to a **whole host of risk factors** (eg, diet, smoking, environmental pollutants, genetics etc). To expect a vitamin pill to work against all these risk factors would seem to be rather unrealistic. With hindsight, therefore, intervention trials evaluating the effect of a vitamin supplement to prevent chronic disease were extremely optimistic.

To put this in context, there is some debate about the **influence of statin drugs to prevent heart attacks, particularly in women.**

Moreover, a careful reading of several of the vitamin trials that have shown either no effect or increased risk of the disease being examined, **shows that the people in the trial often had the disease to start with. For example, in the Heart Outcomes Prevention Evaluation⁷ and the NORVIT⁸ Trial both of which showed no benefit of B vitamins on major cardiovascular events, the people had already got**

cardiovascular disease and some had even had heart attacks. In other words, these were secondary prevention trials.

In summary, vitamins are essential for health and significant numbers of British people have diets lacking in these essential nutrients. Vitamins and minerals should ideally be obtained from the diet, but failing that a supplement can help to fill the nutritional gap. To expect a vitamin pill to prevent a multi-factorial disease which has taken a lifetime to develop, may be unrealistic, particularly if the person already has the disease in question. But what can be expected is that a multivitamin containing a wide variety of vitamins and minerals in the recommended daily amount will ensure that the person's intake does not fall below recommended levels.

-ends-

For further information please contact:

Nicky Smith, Nexus Healthcare

020 7808 9750 / 07867 513 361

Stephanie Lee, Nexus Healthcare

020 78089764

References

1. Rimm EB, Stampfer MJ, Ascherio A, Giovannucci E, Colditz GA, Willett WC. Vitamin E consumption and the risk of coronary heart disease in men. *N Engl J Med* 1993;328(20):1450-6.
2. Holmquist C, Larsson S, Wolk A, de Faire U. Multivitamin supplements are inversely associated with risk of myocardial infarction in men and women--Stockholm Heart Epidemiology Program (SHEEP). *J Nutr* 2003;133(8):2650-4.
3. Homocysteine Lowering Trialists' Collaboration. Dose-dependent effects of folic acid on blood concentrations of homocysteine: a meta-analysis of the randomized trials. *Am J Clin Nutr* 2005;82:806-12.
4. Alfthan G, Aro A, Gey KF. Plasma homocysteine and cardiovascular disease mortality. *Lancet* 1997;349:397.
5. Stephens NG, Parsons A, Schofield PM et al. Randomised controlled trial of vitamin E in patients with coronary disease: Cambridge heart antioxidant study (CHAOS). *Lancet*; 1996; 347:781-6.
6. Toole JF, Malinow MR, Chambless LE et al. Lowering homocysteine in patients with ischemic stroke to prevent recurrent stroke, myocardial infarction, and death: the Vitamin Intervention for Stroke Prevention (VISP) randomized controlled trial. *JAMA* 2004; 291:565-75.
7. Lonn E, Yusuf S, Arnold MJ, Sheridan P, Pogue J, Micks M, et al. Homocysteine lowering with folic acid and B vitamins in vascular disease. *N Engl J Med* 2006;354(15):1567-77.
8. Bonna KH, Njolstad I, Ueland PM, Schirmer H, Tverdal A, Steigen T, et al. Homocysteine lowering and cardiovascular events after acute myocardial infarction. *N Engl J Med* 2006;354(15):1578-88.
9. Giovannucci E, Rimm EB, Ascherio A et al. Alcohol, low methionine, low folate diets and risk of colon cancer in men. *J Natl Cancer Inst* 1995;87:265-73.
10. Giovannucci E, Stampfer MJ, Colditz GA et al. Multivitamin use, folate and colon cancer in women in the nurses' health study. *Ann Intern Med* 1998;129:517-24.
11. Blot WJ, Li JY, Taylor PR. Nutrition intervention trials in Linxian, China: supplementation with specific vitamin/mineral combinations, cancer

- incidence, and disease-specific mortality in the general population. . *J Natl Cancer Inst* 1993;85:1483-92.
12. The Alpha Tocopherol B-CCPSG. The effect of vitamin E and beta carotene on the incidence of lung cancer in male smokers. . *New Engl J Med* 1994;330:1029-35.
 13. Lawson KA, Wright ME, Subar A, Mouw T, Hollenbeck A, Schatzkin A, et al. Multivitamin Use and Risk of Prostate Cancer in the National Institutes of Health-AARP Diet and Health Study. *J. Natl. Cancer Inst.* 2007;99(10):754-764.
 14. Peters U, Foster CB, Chatterjee N, Schatzkin A, Reding D, Andriole GL, et al. Serum selenium and risk of prostate cancer--a nested case-control study. *Am J Clin Nutr* 2007;85(1):209-217.
 15. Henderson L, Irving K, Gregory J, Bates CJ, Prentice A, Parks J, et al. National Diet and Nutrition Survey: adults aged 19 to 64 years. Volume 3. Vitamin and mineral intake and urinary analysis. London: Stationery Office, 2003.
 16. Kiely M, Flynn A, Harrington KE, Robson PJ, O'Connor N, Hannon EM, et al. The efficacy and safety of nutritional supplement use in a representative sample of adults in the North/South Ireland Food Consumption Survey. *Public Health Nutr* 2001;4(5A):1089-97.
 17. Beitz R, Mensink GB, Fischer B, Thamm M. Vitamins--dietary intake and intake from dietary supplements in Germany. *Eur J Clin Nutr* 2002;56(6):539-45.
 18. Schwarzpaul S, Strassburg A, Luhrmann PM, Neuhauser-Berthold M. Intake of vitamin and mineral supplements in an elderly german population. *Ann Nutr Metab* 2006;50(2):155-62.
 19. Archer SL, Stampler J, Moag-Stahlberg A, Van Horn L, Garside D, Chan Q, et al. Association of dietary supplement use with specific micronutrient intakes among middle-aged American men and women: the INTERMAP Study. *J Am Diet Assoc* 2005;105(7):1106-14.
 20. Troppmann L, Gray-Donald K, Johns T. Supplement use: is there any nutritional benefit? *J Am Diet Assoc* 2002;102(6):818-25.
 21. Briefel R, Hanson C, Fox MK, Novak T, Ziegler P. Feeding Infants and Toddlers Study: do vitamin and mineral supplements contribute to nutrient adequacy or excess among US infants and toddlers? *J Am Diet Assoc* 2006;106(1 Suppl 1):S52-65.
 22. Fox MK, Reidy K, Novak T, Ziegler P. Sources of energy and nutrients in the diets of infants and toddlers. *J Am Diet Assoc* 2006;106(1 Suppl 1):S28-42.
 23. Dwyer JT, Garcea AO, Evans M, Li D, Lytle L, Hoelscher D, et al. Do adolescent vitamin-mineral supplement users have better nutrient intakes than nonusers? Observations from the CATCH tracking study. *J Am Diet Assoc* 2001;101(11):1340-6.
 24. Kim SH, Han JH, Keen CL. Vitamin and mineral supplement use by healthy teenagers in Korea: motivating factors and dietary consequences. *Nutrition* 2001;17(5):373-80.
 25. Sichert-Hellert W, Wenz G, Kersting M. Vitamin intakes from supplements and fortified food in German children and adolescents: results from the DONALD study. *J Nutr* 2006;136(5):1329-33.
 26. Stang J, Story MT, Harnack L, Neumark-Sztainer D. Relationships between vitamin and mineral supplement use, dietary intake, and dietary adequacy among adolescents. *J Am Diet Assoc* 2000;100(8):905-10.
 27. Girodon F, Blache D, Monget AL, Lombart M, Brunet-Lecompte P, Arnaud J, et al. Effect of a two-year supplementation with low doses of antioxidant vitamins and/or minerals in elderly subjects on levels of nutrients and antioxidant defense parameters. *J Am Coll Nutr* 1997;16(4):357-65.
 28. Navarro M, Wood RJ. Plasma changes in micronutrients following a multivitamin and mineral supplement in healthy adults. *J Am Coll Nutr* 2003;22(2):124-32.

29. Wolters M, Hermann S, Hahn A. Effects of 6-month multivitamin supplementation on serum concentrations of alpha-tocopherol, beta-carotene, and vitamin C in healthy elderly women. *Int J Vitam Nutr Res* 2004;74(2):161-8.